

Product components

Components	Component number	Size-1	Size-2
		20 RXN	100 RXN
4X ABScript Neo RT Master Mix *	RM21486	100 μ L	500 μ L
20X gDNA Remover Mix	RM21479	20 μ L	100 μ L
Nuclease-free H ₂ O	RM20214	1.25 mL	2 \times 1.25 mL

* 4X ABScript Neo RT Master Mix contains ABScript III Reverse Transcriptase B, RNase Inhibitor, dNTPs, Random Primers/Oligo (dT)₂₀VN Primer Mix, etc.

Product Description

The ABScript Neo RT Master Mix for qPCR with gDNA Remover is a highly efficient and fast cDNA first-strand synthesis master mix, specifically designed for two-step RT-qPCR detection. The 4X ABScript Neo RT Mix in this product contains all the reagents required for the reverse transcription reaction, allowing for a simple and quick reaction setup by adding the RNA template and H₂O. The gDNA Remover Mix in this product can completely remove the genomic DNA remaining in the RNA template and make the qPCR results highly accurate. The dsDNase is heat-sensitive and can be quickly and irreversibly inactivated under high temperature conditions. The dsDNase is heat-sensitive, enabling rapid and irreversible inactivation under high-temperature conditions, requiring only one sample for both genomic DNA removal and reverse transcription reactions in the same tube.

This product is optimized for qPCR, with a proportionally optimized Random Primers/Oligo (dT)₂₀VN Primer Mix that efficiently synthesizes cDNA from all regions of RNA transcription, ensuring maximum authenticity and repeatability of qPCR results. Reverse transcription products generated using this kit are compatible with SYBR Green and probe qPCR and can be used in combination with corresponding reagents, tailored to specific experimental purposes, for high-performance gene expression analysis.

Storage

-20°C

Precautions for Use

1. Please briefly centrifuge them to the bottom of the tube before use, and gently pipette to mix prior to use.
2. Random Primer and Oligo (dT)₂₀VN Primer have been added to this product, thus gene-specific primers cannot be used.
3. The reverse transcription product (cDNA) obtained by using this product is only suitable for qPCR reaction and is not suitable for long fragment PCR amplification in downstream experiments such as cloning. If necessary, you can use ABScript II cDNA First-Strand Synthesis Kit (ABclonal RK20400) to conduct experiments.
4. Replace pipette tips when transferring reagents to avoid cross-contamination.

Requirements

1. Materials and Equipment: 1.5 mL RNase-free EP tubes, 200 μ L RNase-free PCR tubes, RNase-free pipette tips, pipettors, PCR instrument (and qPCR instrument), ice or ice box.
2. RNA: Complete and high-quality RNA is essential for obtaining high quality cDNA.
3. Ensure RNA is not degraded or contaminated before the experiment. If RNA contains a complex secondary structure or a high GC content, it can be incubated at 65°C for 5 minutes (and immediately on ice) before reverse transcription.

Experimental procedure

1. Reverse transcription
 - (1) Reverse transcription reaction system

Add the components to the RNase-free PCR tube on ice according to the following recommendations, mix well and centrifuge briefly.

Components	Volume
4X ABScript Neo RT Master Mix	5 μ L
20X gDNA Remover Mix	1 μ L
Total RNA	10 pg -1 μ g *
Nuclease-free H ₂ O	to 20 μ L

* Add the appropriate amount of RNA according to the experimental requirements. When the RNA template is too much, make sure that the RNA is soluble in water and not in TE Buffer, as TE inhibits the reverse transcription reaction.

(2) Reverse transcription reaction procedure

Temperature	Time
37 °C	2 min
55 °C	15 min
85 °C	5 min
4 °C	Hold

* Product can be applied immediately to the subsequent qPCR reaction, or stored at -20°C . Avoid repeated freezing and thawing.

2. qPCR

The following is after used this product for reverse transcription, select 2X Universal SYBR Green Fast qPCR Mix (ABclonal RK21203) reagent to carry out qPCR reaction in StepOnePlus Real-Time PCR System.

* Please read the instrument operation manual before the experiment.

(1) qPCR reaction system (Take 20 μ L as an example)

Component	Volume
2X Universal SYBR Green Fast qPCR Mix	10 μ L
cDNA product (RT reaction liquid)	X μ L*
Forward Primer (10 μ M)	0.4 μ L
Reverse Primer (10 μ M)	0.4 μ L
Nuclease-free H ₂ O	to 20 μ L

* It is suggested that the volume of the template does not exceed the 1/10 volume of the qPCR reaction, or the Nuclease-free H₂O is used to dilute the cDNA product (RT reaction liquid) and then add to the reaction system.

(2) qPCR reaction procedure (two-step)

Step	Temperature	Time	Cycles
Stage1	95 °C	3 min	1 cycle
Stage2	95 °C	5 sec	40 cycles
	60 °C	30 sec	

Melt Curve (automatic instrument setting)

Analysis of result

The amplification curve and melting curve of qPCR were confirmed after the reaction, and then the standard curve was made for quantitative analysis. The method of analysis is referred to the manual of the instrument operation.