

ABScript II Reverse Transcriptase



WEB: www.abclonal.com

Catalog: RK21400

Size: 4,000 U / 10,000 U

Concentration: 200,000 U/ml

Components:

ABScript II Reverse Transcriptase (200,000 U/ml)	RM21400
5X First-Strand Buffer	RM20109
100 mM DTT (10X)	RM20117

Storage Conditions:

20 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.01% IGEPAL® CA-630, pH 7.5 @ 25 °C

Heat Inactivation: 65 °C for 20 min

Product Description

ABScript II Reverse Transcriptase is a recombinant M-MuLV reverse transcriptase with reduced RNase H activity and increased thermostability. It can be used to synthesize first-strand cDNA at higher temperatures than the wild type M-MuLV. The enzyme is active up to 48 °C, providing higher specificity, higher yield of cDNA and more full-length cDNA product up to 12 kb.

Product Source: The gene encoding a mutant M-MuLV Reverse Transcriptase (RNase H) is expressed in *E. coli* and purified to near homogeneity.

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 1 nmol of dTTP into acid-insoluble material in 10 minutes at 37 °C using poly(rA)•oligo(dT)₁₈ as template, with a total reaction volume of 50 µl.

Reaction Conditions:

1X First-Strand Reaction Buffer, 10 mM DTT, 200 units ABScript II Reverse Transcriptase, supplemented with 0.5 mM dNTPs (not included) and 5 µM dT₂₃VN (not included). Incubate at 42 °C for 50 minutes. If random primers are used, a 10-minute incubation at room temperature is recommended before transferring to 42 °C.

1X First-Strand Reaction Buffer:

50 mM Tris-HCl, 75 mM KCl, 3 mM MgCl₂, pH 8.3 @ 25 °C

Storage Temperature: -20 °C

Instructions

➤ **First Strand cDNA Synthesis (Quick Protocol)**

Thaw components on ice and mix by inverting several times.

1. Mix the following components and incubate at 42 °C** for 1 hour. If Random Primer Mix is used, an incubation step at 25 °C for 5 minutes is recommended before the 42 °C incubation.

COMPONENT	VOLUME
Nuclease-free H ₂ O	Up to 20 µl
5X First-Strand Buffer	4 µl
d(T) ₂₃ VN (50 µM) or Random Primer Mix (60 µM)	2 µl
100 mM DTT	2 µl
RNase Inhibitor (40 U/µl)	0.2 µl
Template RNA	up to 1 µg*
10 mM dNTP	1 µl
ABScript II RT (200 U/µl)	1 µl

* 1 ng-1 µg total RNA or 50 pg-100 ng poly(A)-RNA

** ABScript II Reverse Transcriptase can be used at 42–48 °C.

2. Inactivate the enzyme at 65 °C for 20 minutes. For downstream PCR application, the volume of cDNA product should not exceed 1/10 of the PCR reaction volume.

➤ **First Strand cDNA Synthesis (Standard Protocol)**

If denaturation of template RNA is desired, use the following protocol.

1. Mix RNA sample and primer d(T)₂₃VN in a sterile RNase-free microfuge tube.

COMPONENT	VOLUME
Total RNA	up to 1 µg*
d(T) ₂₃ VN (50 µM) or Random Primer Mix (60 µM)	2 µl
10 mM dNTP	1 µl
Nuclease-free H ₂ O	Up To 10 µl

- Denature sample RNA/primer for 5 minutes at 65 °C. Spin briefly and put on ice immediately.
- Add the following components to the tube

COMPONENT	VOLUME
Nuclease-free H ₂ O	Two Steps Total Up to 20 µl
5X First-Strand Buffer	4 µl
100 mM DTT	2 µl
RNase Inhibitor (40 U/µl)	0.2 µl
ABScript II RT (200 U/µl)	1 µl

- Incubate the 20 µl cDNA synthesis reaction at 42 °C** for one hour. If Random Primer Mix is used, an incubation step at 25 °C for 5 minutes is recommended before the 42 °C incubation.
- Inactivate the enzyme at 65 °C for 20 minutes. The cDNA product should be stored at -20 °C. In general, the volume of cDNA product should not exceed 1/10 of the PCR reaction volume.

* 1 ng-1 µg total RNA or 50 pg-100 ng poly(A)-RNA

** ABScript II Reverse Transcriptase can be used at 42–48 °C.

➤ First strand cDNA Synthesis (No-RT Negative Control Reaction)

- Mix the following components and incubate at 42 °C for 1 hour:

COMPONENT	VOLUME
Template RNA	up to 1 µg*
d(T) ₂₃ VN (50 µM) or Random Primer Mix (60 µM)	2 µl
5X First-Strand Buffer	4 µl
100 mM DTT	2 µl
10 mM dNTP Mix	1 µl
RNase Inhibitor (40 U/µl)	0.2 µl
Nuclease-free H ₂ O	Up to 20 µl

* 1 ng-1 µg total RNA or 50 pg-100 ng poly(A)-RNA

QC Process:

- ◆ Purity is above 95% detected by SDS-PAGE.
- ◆ No exonuclease, nuclease, RNase contamination.
- ◆ Host genomic DNA is no residual detected by PCR.

Additional Products:

RNase Inhibitor (40,000 U/mL) is available separately (Cat. NO. RK20401).