

HS *Taq* DNA Polymerase M101

(5,000 U/mL)

Version: 16G19v1.1

Catalog: RK26201

Size: 250 U / 1,000 U

Components:

| | |
|---|---------|
| HS <i>Taq</i> DNA Polymerase M101 (5,000 U/mL) | RM29201 |
| 10X PCR Reaction Buffer, Mg ²⁺ plus | RM20101 |

Product Description

HS *Taq* DNA Polymerase M101 is an antibody-blocked thermostable *Taq* DNA polymerase with a theoretical molecular weight of 94 KD. It has 5' → 3' polymerase activity and 5' → 3' exonuclease activity without 3' → 5' exonuclease activity. The PCR product has A-tailing at the 3' end, which can be cloned with dT/dU vector.

The polymerase activity of this product is completely inhibited at room temperature, avoiding non-specific amplification and primer dimer during the preparation of PCR reaction system. During the pre-denaturation process at 95 °C, the antibody will denature and be inactivated, releasing the activity of *Taq* DNA polymerase. This step will not affect the subsequent PCR reaction, but also increase the specificity of PCR. This product is applicable to qPCR and RT-qPCR.

Product Components

| Component | 250 U | 1,000 U |
|---|---------|---------|
| HS <i>Taq</i> DNA Polymerase M101 (5,000 U/mL) | 50 µL | 200 µL |
| 10X PCR Reaction Buffer, Mg ²⁺ plus* | 1.25 mL | 1.25 mL |

*, Note: 10X PCR Reaction Buffer, Mg²⁺ plus should only be used as reaction buffer for HS *Taq* DNA Polymerase M101.

Product Source

This product is expressed in Escherichia coli cloned with Thermus aquaticus YT-1 *Taq* DNA Polymerase gene and

is purified through multiple steps.

Unit Definition

One unit (U) is defined as the amount of enzyme that incorporates 10 nM of dNTPs into acid-insoluble products in 30 min at 75°C.

1X PCR Reaction Buffer, Mg²⁺ plus Components

20 mM Tris-HCl, 10 mM (NH₄)₂SO₄, 10 mM KCl, 2 mM MgSO₄, 0.1% Triton X-100, pH 8.8 @ 25°C

Storage

Store at -20°C.

Storage Solution for Enzymes

10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5% Tween 20, 0.5% NP-40, 50% Glycerol, pH 7.4@ 25°C

Recommendations

Recommended PCR reaction

PCR reaction system (50 µL) *

| Components | Volume (50 µL reaction) |
|--------------------------------------|----------------------------|
| ddH ₂ O | to 50 µL |
| 10X PCR Buffer | 5 µL |
| 10 mM dNTPs | 1 µL |
| Upstream Primer (10 µM) | 1 µL |
| Downstream Primer (10 µM) | 1 µL |
| Template DNA | Variable |
| HS <i>Taq</i> DNA Polymerase M101 ** | 5 U |

*, Note: Gently mix the reaction system, and if necessary, collect the reagent to the bottom of the tube by rapid centrifugation. If the PCR instrument without hot lid is used, the surface of the reaction system can be covered with a layer of mineral oil to prevent the evaporation of the solution.

******, Note: The amount of HS Taq DNA Polymerase M101 can be adjusted with the range of 1-25 U according to the situation in the 50 μ L reaction system.

Reaction Process

| Step | Temperature | Time | Cycles |
|----------------------|-------------|----------|--------|
| Initial Denaturation | 95°C | 5 min | 1 |
| Denaturation | 95°C | 15-30 s | 25-40* |
| Annealing | 45-68°C | 15-60 s | |
| Extension | 68°C** | 60 s/kb | |
| Final Extension | 68°C | 5 min | 1 |
| Hold | 4-10°C | ∞ | 1 |

*****, Note: Generally, sufficient PCR products can be obtained after 25-40 cycles. If it is necessary to detect low copy genes, the number of cycles can be increased to 45.

******, Note: The extension temperature of 68 °C is recommended. The extension time is related to the length of the amplified fragment. The amplification time can be calculated according to the amplification speed of 60 s/ KB; After the PCR cycle, it needs to be extended at 68 °C for another 5 minutes.